



# USF Breast Health

## MEDICATIONS THAT NEED TO BE STOPPED PRIOR TO SURGERY:

The following medications all contain aspirin or aspirin-like products and will contribute to “thinning your blood” and putting you at risk for bleeding during surgery. If you are taking any of these products including Vitamin E, our recommendation is that you stop these 7 days prior to surgery. You may take Tylenol, (acetaminophen) during this time. **EPHEDRINE AND PHENDIMETRAZINE: you MUST be off these medications for 14 days before surgery. If you are currently on a Coumadin product, please discuss with your doctor.**

- DIET PILLS OF ANY TYPE MUST BE STOPPED 14 DAYS PRIOR TO SURGERY!
- Advil
- Acuprin
- Alka-Seltzer products
- Anaprox
- Ansaid
- Anacin products
- Ascriptin with Codeine
- Arthritis Strength Bufferin
- Aspirin (all forms and brands)
- ASA/ASA enesals
- Ascriptin
- Aspergum
- Clinoril
- Coumadin
- Darvon Compound
- Daypro
- Disalcid
- Doan pills
- Dolobid
- Ecotrin products
- Empirin
- Empirin with Codeine
- Equagesic
- Excedrin Products
- Feldene
- Fiorinol with Codeine
- Fish Oil pills
- Flexiril
- Goody's
- Herbal tea of any kind
- Ibuprofen products
- Lortab ASA
- Medipren
- Methocarbamol with ASDA
- Midol
- Mobigesic
- Momentum
- Motrin
- Nalfon
- Nardit
- Naprosyn
- Naproxen
- Norgesic
- Nuprin
- Pamprin
- Peto Bismol
- Percodan
- Propoxyphene Compound
- Rufen
- Toradol
- Tolectin
- Tometin
- Trilasate
- Vanquish
- Vioxx
- Vitamin E
- Voltaren
- Zorprin
- Zylprim

