



Complete for 20 systematically selected postpartum (PP) women (sampling method on the back) admitted to your hospital for delivery regardless of infant outcome

		STUDY ID #			
DEMOGRAPHICS					
Delivery Month Year	Saturday/Sunday/Holiday discharge ☐ Yes ☐	turday/Sunday/Holiday discharge 🗆 Yes 🗆 No 🛮 Maternal ag		_	
☐ Medicaid/Medicaid plans Type of ☐ Private insurance ☐ Self-pay ☐ Other ☐ Unknown	Mother's	Mother's Ethnicity	☐ Hispanic Mother's ☐ Haitian Ethnicity ☐ Non-Hispanic/Non-Haitian ☐ Unknown		
Prenatal care started in □ I / II trimester □ III trimester □ No prenatal care □ Unknown	Mother's Preferred Language English Unknown Spanish Creole Other:	Route of Delivery	□ Vaginal □ Cesarean		
POSTPARTUM CARE					
			Yes	No	
Was a Maternal Discharge Risk Assessment performed?					
Was the patient (pt.) verbally instructed on the benefits of early risk-appropriate PP visits/Post-Birth Health Check and given written materials?					
Was the pt. verbally instructed about PP Warning Signs and given PP warning signs written materials?					
Was the pt. verbally instructed about the benefits of and options for pregnancy spacing, family planning, and contraceptive choice and given written materials?					
Was a PP Discharge Assessment (vital signs and response) conducted just prior to discharge?					
POSTPARTUM VISITS					
How many days after delivery were Postpartum Visits scheduled prior to discharge (check all that apply)? □ <7 days □ 7-14 days □ 15 - 21 days □ > 21 days □ Not scheduled/mother instru					
PP High risk? ☐ Yes → check condition(s) below ☐ No Referrals scheduled and medicati					
- Chi offic i i i v, gestational i i i v, pre eclampsia, eclampsia, maternal near t alsease, or		l prior to dis k all that ap			
☐ Hx of venous thromboembolism (DVT or pulmonary embolism) /on anticoagulation Specialty appoin		ntment			
☐ C-section or 3rd or 4th degree vaginal laceration Mental/Behavio appointment			ral Health		
☐ Positive screen for Substance Use Disorder Healthy Start/ho			ome visiting		
☐ Feeling unsafe at home / Positive for Intimate Partner Violence Medicaid Case M			/Janager		
☐ Positive Edinburgh Postnatal Depression Scale Hospital financia			al counselor		
☐ Requested/required additional community resources Appropriate me			dication		
☐ Other Naloxone kit/Rx					



SAMPLING

- If your hospital has 40 births per month or more: report 20 systematically selected discharged deliveries per month. Selection process: Start by dividing the total number of delivery discharges that occurred at your facility in a given month by 20. Then select every nth chart where n is the result of that division. e.g. Your hospital had 105 discharged deliveries in June. Divide 105 by 20. 5 is your nth for June. Report data on every 5th chart.
- If your hospital has less than 40 births per month: report the first 10 delivery charts per month.

DATA DEFINITIONS

Maternal age: the age of the mother at the time of delivery. Completed years only. Do not round up.

PP Discharge Assessment: check blood pressure, heart rate, respiratory rate, <u>and</u> body temperature **just prior to discharge -** refer to FPQC's PP Discharge Assessment for appropriate response.

Post-Birth Health Check: patient preferred language for the early risk-appropriate PP visit within 2 weeks postpartum

MATERNAL DISCHARGE RISK ASSESSMENT

Guidelines for postpartum management are to be set by your hospital. Patients should be assessed for the following at minumum:

Chronic hypertension, gestational hypertension, pre- eclampsia, eclampsia or maternal heart disease, or related conditions	Schedule blood pressure check in 2-3 days and appointment with OB or PCP in 1-2 weeks. If yes to maternal heart disease, schedule appointment with cardiology in 1-2 weeks.		
History of venous thromboembolism (DVT or pulmonary embolism) this pregnancy or on anticoagulation prior to delivery	Ensure patient has 6 weeks of medication for anticoagulation in hand prior to discharge.		
C-section or 3rd or 4th degree vaginal laceration	Schedule for 1–2-week incision check with OB.		
Positive screen for Substance Use Disorder	Perform SBIRT; refer for MAT/MOUD, Narcan kit, OB follow up in 1-2 weeks		
Feeling unsafe at home or positive screen for Intimate Partner Violence	Refer to case manager or social worker for assessment prior to discharge.		
Feelings of depression, hopelessness or disinterest	Screen with Edinburgh Postnatal Depression Scale, contact OB provider, and schedule follow up for mood check in 1-2 weeks. Consider psych consult		
Need for additional community resources	Consult social worker, refer to Healthy Start, Medicaid Case Manager, or hospital financial counselor.		